

The comparison of the effect of acute moderate and high-intensity exercise on the uncoupling protein -1 secretion

Desiana Merawati^{1abe}, Sugiharto^{1abe}, Adi Pranoto^{2cdf}, Olivia Andiana^{1bc}, Prayogi Dwina Angga^{3bc}

¹Department of Sport Sciences, Faculty of Sport Sciences, Universitas Negeri Malang, Indonesia

²Doctoral Program of Medical Sciences, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

³Department of Elementary Education, Faculty of Teacher Training and Education Universitas Mataram, Indonesia

Authors contribution: a – Preparing concepts; b – Formulating methods; c – Conducting research; d – Processing results; e – Interpretation and conclusions; f - Editing the final version

Received: 5 Maret 2022; Revised: 29 June 2022; Accepted: 18 August 2022

Abstract

Physical activities accelerate the secretion of the uncoupling protein-1 (UCP-1), which reduces the risk of metabolic disorders and the prevalence of obesity. This research aimed to investigate the effects of acute physical activities on UCP-1 secretion. A total of 20 males (aged between 19-20 years old) were divided into groups with either a 30-minute moderate-intensity exercise (MIE) or high-intensity exercise (HIE). The UCP-1 expression was measured using Enzyme-Linked Immunosorbent Assay (ELISA) (pre and 5 minutes post the exercise). The pre and post-UCP-1 data in each group were compared with a paired t-test, while pre and post UCP data between groups were analyzed using the Independent-Sample T-test. The pre and post test UCP-1 in the MIE was (4.16 ± 0.89) ng/mL and (4.55 ± 1.36) ng/mL ($P > 0.05$), while the pre and post test UCP-1 in the HIE were (4.09 ± 0.53) ng/mL and (5.06 ± 1.02) ($P < 0.05$). No significant difference in pretest UCP-1, however was found between groups ($P > 0.05$) and posted UCP-1 ($P > 0.05$). In conclusion, 30 minutes of high-intensity exercise is required to increase UCP-1 in young adults significantly.

Keywords: moderate-intensity, high-intensity, exercise, UCP-1

INTRODUCTION

Currently, obesity is perceived as a world pandemic that has affected 600 million adults worldwide (Rodrigues et al., 2018). WHO survey estimates that 10.8 % of men and 14.9% of women encounter obesity with a body mass index ≥ 30 (Appari et al., 2018). The fact obesity belongs to 3 out of 5 high-risk diseases (Kim et al., 2018). The expected obesity prevalence in the Asia Pacific is 11.9%-37.1%, foreseen as the trigger of accelerating cardiovascular diseases (CVD) and type 2 diabetes mellitus

Correspondence author: Desiana Merawati, Universitas Negeri Semarang, Indonesia.
Email: desiana.merawati.fik@um.ac.id



Jurnal SPORTIF: Jurnal Penelitian Pembelajaran is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/)

prevalence ([Herningtyas and Ng, 2019](#)), as well as the prevalence of high blood pressure, dyslipidemia, and at risk of complication ([Kim et al., 2018](#)). Besides, it possibly escalates morbidity and premature death ([Flouris et al., 2017](#)), and threatens people's economic and social health ([Appari et al., 2018](#)). Physical activation increases aerobic metabolism and browning activity ([Purdom et al., 2018](#)) thus, it enables increasing UCP-1 secretion ([Kim and Plutzky, 2016](#)). The body's response and adaptation to exercise activities are complex. Therefore, suitable and sufficient physical exercises to improve UCP-1 expression is still required.

Exercise is a healthy and fun activity that includes all systems ([Huh, 2018](#)). In addition, exercise can also increase fat oxidation and the ability of skeletal muscles to utilize fat ([Jabbour and Iancu, 2017](#)). However, an exercise that is done inappropriately has the potential to trigger metabolic disorders ([Brondani et al., 2012](#)). Metabolic disorders can result in impaired fat oxidation and reduce protein expression in mitochondrial membranes ([Huh, 2018](#)). Yet, the right exercise intensity boosts aerobic metabolism and browning activity ([Jabbour and Iancu, 2017](#)). Recent research reports that low-moderate physical activities escalate lipid metabolism, but heavy intensity exercises tend to decrease lipid metabolism ([Oh et al., 2017](#)). In conclusion, exercise intensity and duration correlate with fat metabolism transforming protein composition in mitochondria and causing an efficient oxidation process by altering H proton going through the membrane to transport electrons used for managing ATP synthesis ([Befroy et al., 2008](#)). The increase in lipid metabolism is initiated by the acceleration of irisin secretion produced by escalated muscle contraction during physical activities ([Brondani et al., 2012](#)).

Expansion of irisin drives increasing browning ([Oh et al., 2017](#)) activating UCP-1 expression so that the thermogenesis is improved ([Brondani et al., 2012](#)). However, not all physical activities accelerate UCP-1. It is reported that 12 weeks of endurance exercises in healthy and prediabetic adults subject gives no significant effect on fat oxidation ([Zhang et al., 2017](#)). Besides, physical activities do not affect white adipose tissue

(Dinas et al., 2017). UCP-1 is located inside the mitochondrial membrane of brownish adipose. It acts as a transmembrane protein that forces ATP synthesis through fat oxidation and thermogenesis that relies on exercise (Fenzl and Kiefer, 2014). Continual moderate and high-intensity physical activities increase UCP-1 expression and gene expression from fatty acid transportation (Khalafi et al., 2020a). However, the 60%-64% VO₂max intensity exercise on the trained person and 47%-52% VO₂max intensity exercise on the untrained person extend fatty oxidation (Oh et al., 2017).

In contrast, running with 100% aerobic speed better repairs fatty oxidation and body composition than moderate intensity (Jabbour and Iancu, 2017). Other researchers state that a long duration of moderate exercise is perceived as effective for fatty oxidation (Purdom et al., 2018). Yet, research also reveals that six weeks of moderate-intensity exercise produces no difference in UCP-1 expression activity (Dinas et al., 2017). In other words, fatty oxidation is not only affected by intensity, duration and types of physical activities but also the training, obesity level, types of obesity (visceral or subcutaneous fat), food supply (before or after meal) and the food composition (fat percentage), as well as the genetic factors (Oh et al., 2017). Chronic and acute physical activities activate brown fat (Gorski et al., 2018) by increasing PGC-1 α on skeletal muscle (Dinas et al., 2017). Therefore, this study examines the effect of high and moderate physical activities on the expression of UCP-1 in young adults to decrease and prevent the accelerating obesity prevalence. The subjects used in this study have criteria, namely male, aged 19-20 years, have a good level of physical fitness, the subject is a trained person but not an amateur or professional athlete.

METHODS

Study design

This study uses a true experimental method with a randomized pretest-posttest group design. A total of 20 19-20 years old young adults, body mass index (BMI) of 18.5-24.9 kg/m², normal blood pressure, regular resting heart rate (RHR), excellent physical freshness level, and healthy

hemoglobin (Hb) contribute to studies. They were randomly divided into two groups: AMIP (n=10) and AHIP (n=10). The Ethical Committee approved all procedures of the study of the Faculty of Medicine, Brawijaya University, Malang, with registration number 106/EC/KEPK/04/2018.

Physical activity protocol

The exercise was completed by running on a treadmill for 40 minutes (Rejeki et al., 2021; Rejeki et al., 2022; Andarianto et al., 2022) with a high-intensity (75-85% HRmax) and moderate-intensity (65-75% HRmax). Intensity measurement was carried out using a measurement formulation of HRmax: HRmax – age in years (220 – age in years) (Sugiharto et al., 2021; Tsuchiya et al., 2014). During that activity, the subjects' heartbeat rates were monitored by the polar heart rate monitor.

Data collection

The assessment of participants' anthropometry, physiology and physical fitness involved body height, body weight, body mass index (BMI) and blood pressure, Hb and physical fitness measurement with a multi-stage fitness test (MFT) 20 M (Paradisis et al., 2014; Sugiharto et al., 2022).

In addition, the blood draw for UCP-1 expression analysis from the cubital vein (Daskalopoulou et al., 2014) was performed during the before-exercise dan 5 min after exercise (Huh et al., 2014). After that, the blood was centrifuged. The serum was divided and saved at -80°C temperature (Tsuchiya et al., 2015). The UCP-1 secretion measurement was accomplished using ELISA (Catalog No. E-EL-H1661; Elabscience, Inc., China) (Adji et al., 2021; Sugiharto et al., 2021).

Statistical analysis

Data were analyzed using SPSS version 17 (SPSS Inc., Chicago, IL, USA). The data was first tested using a descriptive test, then the data was tested for normality using the Shapiro Wilk test. Furthermore, in assessing the differences between those moderate and high-intensity exercises between pre or post-test using Paired Samples T-test, with a significance level of ($P < 0.05$), meanwhile to assess differences between groups, the

Independent-Samples T-test was performed, with a significance level of ($P < 0.05$).

Research procedure

The time it takes the issue to provide treatment, i.e. one intervention with high and moderate intensity treadmill exercise for 40 minutes. The technique used for taking the subject is using the Consecutive Sampling technique. The division of groups is done by matching and then using a zig-zag system. Data collection is done by checking the level of UCP-1 between before and 5 minutes after the intervention using the ELISA Kit. The treatment was observed by monitoring heart rate during exercise using a POLAR heart rate monitor H10.

RESULTS

The descriptive analysis of each participant's body profile is presented in Table 1.

Table 1. Characteristics of the Subjects

Variable	n	Group		p-value
		MIE (n=10)	HIE (n=10)	
Age (years)	10	20.40±0.69	20.20±0.79	0.145
Body Height (m)	10	1.65±0.05	1.67±0.04	0.586
Body Weight (kg)	10	60.20±5.18	59.50±6.48	0.570
BMI (kg/m ²)	10	22.06±1.52	21.25±1.40	0.092
SBP (mmHg)	10	118.00±4.22	114.00±8.43	0.254
DBP (mmHg)	10	81.00±3.16	77.00±4.83	0.120
RHR (bpm)	10	72.40±10.06	70.80±8.44	0.911
VO _{2max} (mL/kg/min)	10	42.65±2.72	42.78±3.08	0.992
Hb (g/dL)	10	15.38±1.33	16.40±2.41	0.625

Independent-Sample T-Test. Data are presented as mean±SD. BMI: Body mass index; SBP: Systolic blood pressure; DBP: Diastolic blood pressure; RHR: Resting heart rate; VO_{2max}: Maximum oxygen volume; Hb: Hemoglobin; MIE: moderate-intensity exercise; HIE: high-intensity exercise.

The Independent-Sample T-test presented in Table 1 displays no essential differences among the healthy subjects' characteristics in both groups. Additionally, the results of the UCP-1 secretion analysis between before and 5 min after exercise in each group are shown in Figure 1.




Figure 1. Graphic of average UCP-1 expression analysis between MIE and HIE

Based on the results of the normality test in Figures 1 and 2 show that the data has a normal distribution ($P > 0.05$). The Independent-Sample T-test in Figure 1 exhibits no significantly different average UCP-1 secretion between MIE vs. HIE on pre ($P > 0.05$) and post-test ($P > 0.05$). The results of the analysis of UCP-1 secretion before and 5 min after exercise in each group are shown in Figure 2. The Paired Samples T-test presented in Figure 2 displays no essential average UCP-1 secretion between before and 5 min after exercise on the AMIP group ($P > 0.05$). In contrast, there is a significant difference in UCP-1 secretion before and 5 min after exercise on AHIP ($P < 0.05$).




Figure 2. Mean of UCP-1 secretion analysis between before exercise and 5 min after exercise. (*) Significant with $P < 0.05$ by Paired Samples T-test.

DISCUSSION

Exercise is one of the strategies and no-drug approaches to prevent the accelerating chronic disease prevalence, such as obesity, type 2 diabetes mellitus, and metabolic diseases (Dewal & Stanford, 2018). Therefore, exercise should be a lifestyle to prevent increasing chronic disease prevalence (Müller et al., 2016), and regulate energy balance and body function (Garneau et al., 2020). Chronic diseases and premature death will decrease significantly (Leal et al., 2018). A molecular physiology study on exercise is required to counter the rise of chronic diseases. Therefore, this research aims to analyze UCP-1 expression after high- and moderate-intensity acute exercise in young adults. The results demonstrate no difference in UCP-1 expression in both physical activities ($P>0.05$). Thirty minutes of high- and moderate-intensity acute exercise on young adults with excellent freshness levels have a similar response toward UCP-1 expression. A previous study by Kim et al. (2018), giving 30 minutes of different exercise intensity on the treadmill to lab animals, also reveals no significant difference in UCP-1 expression.

In addition, research using obese white Wistar rat lab animal given high-intensity exercise on a treadmill with 85%-90% maximum speed, along with continuous moderate-intensity with 65%-70% maximum speed toward Adipogenesis and Browning on adipose tissue reports no significant increase of UCP-1 on both groups (Khalafi et al., 2020b). Previous research also finds that exercise enhances UCP-1 expression and sympathetic nerve stimulation (Dewal and Stanford, 2018). In addition, the results of this study also revealed accelerated UCP-1 secretion occurred in both groups. The develop UCP-1 is mediated by the raise of gen 1-alpha (PGC-1 α) periferisome proliferator-activated gen receptor activation on human skeletal muscle triggered by exercise (Reisi et al., 2016). PGC-1 α is a co-transcription regulator that promotes various transcription managing complex gene regulatory networks that engage in the mitochondrial tissue content control producing the brown adipose tissue (BAT) production, while

PGC-1 α improves protein with five fibronectin type III 5 (FNDC5) domain that is divided and released to blood flow as irisin (Dinas et al., 2017).

The results of this research present that high-intensity exercise produces more extensive UCP-1 expression than moderate-intensity activity. The exercise intensity affects the increase of muscle contraction and accelerates sympathetic nerve stimulation (Aldiss et al., 2018). The intensively developed muscle construction during the exercise drives the metabolism escalation that inspires the advance of PGC-1 α , which is higher on the high-intensity than the moderate-intensity (Shirvani and Arabzadeh, 2020). PGC1- α is the mediator of energy metabolism programming. It controls mitochondrial biogenesis and oxidative metabolism, such as cell types, that stimulate UCP-1 expression (Dinas et al., 2017), in chronic and acute physical activities (Gorski et al., 2018). The accelerated UCP-1 expression comes from physical activity provocation mediated by co-activator PGC1 α , increasing FNDC5 in muscle, secreted irisin, that boosts thermogenesis gene stimulant on adipocytes. Irisin functions as the signal administrator of energy secretion from the muscle directly communicated with adipose tissue and causes the browning, accelerating the metabolic profile of white adipose tissue (WAT) to improve the whole body energy release (Qiu et al., 2018).

In addition, adipose tissue is a dynamic tissue for energy metabolism, while brown adipose tissue operates as active tissue metabolically contributes toward thermogenesis through activating UCP-1 (Khalafi et al., 2020b). On the other hand, thermogenetic is commanded by the hypothalamus. Thus, body temperature shift leads to thermosensory stimulating dorsomedial hypothalamic as an adaptive thermogenesis control mediated by the hypothalamus (Morrison 2016). A study of rats exposed to cold reports that thermosensory input enforced through the preoptic median prompts the dorsomedial hypothalamus (DMH) so that the sympathetic nerve activity is increased (Fuller-Jackson and Henry, 2018).

The increased catecholamine secretion may affect the escalated UCP-1 expression during the high-intensity exercise (Sanchez-Delgado et al.,

2015). The adrenergic receptor captures the extension of catecholamine secretion, increasing cyclic adenosine monophosphate and then activating protein kinase A, drives lipolysis and UCP-1 secretion (Rodrigues et al., 2018). Noradrenaline binds β_3 , α_1 , and α_2 -adrenergic receptors. It generates the upsurge signal production that disturbs the process of lipolysis, thermogenesis, and gene expression. This way, β_3 on BAT improves metabolism that activates UCP-1. UCP-1 is a non-related protein in mitochondria responsible for proton translocation through the respiratory chain producing heat. Besides, a continuous β -adrenergic stimulant from BAT produces thermogenic activity that develops UCP-1 expression and BAT hyperplasia accountable for energy balance maintenance (de Queiroz et al., 2012) and decreasing reactive oxygen species (ROS) (Brondani et al., 2012).

A more intensive skeletal muscle construction influences the high degree of UCP-1 in high-intensity activity than in moderate-intensity exercise (Huh, 2018). The muscle construction intensity affects the increase of energy necessity, oxygen uptake, and more active AMPK, improving the active PGC-1 α and the developed thermogenesis (Khalafi et al., 2020b). Irisin discharge begins the expansion of mitochondrial content and UCP-1 expression within white adipose tissue (WAT), which improves thermogenic (Mai et al., 2020). Energy expenditure also induces PGC-1 α transcription muscle regulation, stimulates the FNDC5 protein-membrane increase, improves iron discharge, creates biogenesis in mitochondria, and UCP-1 expression, accelerates heat production, and increases energy expenditure (Moienneia and Attarzadeh Hosseini, 2016). It is proven that the energy expenditure profile during high-intensity exercise is more comprehensive than moderate-intensity, even without an essential different temperature increase (Sugiharto et al., 2019a).

Additionally, the developed muscle construction intensity on high-intensity exercise accelerates higher energy requirement and oxygen need, creating the surge of sympathetic nerve stimulant to free catecholamines, affecting UCP-1 expression that browning in fulfilling the energy demand

(Dewal and Stanford, 2018). Previous research reports higher heart rate and blood pressure escalation in high-intensity than moderate-intensity exercise (Sugiharto et al., 2019b). Intense muscle contraction during high-intensity exercise is one of the factors that affect sympathetic nerve stimulus and catecholamines secretion that binds the β -adrenergic receptor, along with G protein; it drives the increase of adenylate cyclase activity, AMP cyclic (cAMP) activation, A-kinase protein, and p38MAPK, stimulates lipolysis that activates UCP-1 (Sanchez-Delgado et al., 2015). Research results reveal that intense skeletal muscle contraction during high-intensity exercise is the primary determinant of different UCP-1 expressions, even when the UCP-1 secretion in the second group is not critically different. There is a possibility that the exercise duration, types of muscle contraction, and excellent subjects' freshness level have not significantly accelerated UCP-1 secretion. The exercise duration provides an authentic contribution toward UCP-1 expression. The research presents that 30 minutes of exercise with 60%-70% VO₂max intensity provides no significant UCP-1 expression (Kim et al., 2018). Therefore, future research should pay attention to exercise duration, different activity modifications, and additional browning markers related to UCP-1 levels, such as IL-6.

CONCLUSION

Exercise. The results of the study showed no difference in the expression of UCP-1 between pre and post-test in MIE. In contrast, the UCP-1 of pre and post-test on AHIP exhibits significant differences. The 30 minutes acute high-intensity exercise substantially accelerates the expression of UCP-1 than a severe moderate-intensity exercise in young adults. Thus, high-intensity exercise is proposed to improve UCP-1 expression for young adults and trained groups by still observing the level of training, heart rate, and blood pressure during the exercise. In contrast, untrained people are not suggested to use the high-intensity exercise. Besides, future researchers are expected to analyze untrained or obesity groups with different activity models.

CONFLICT OF INTEREST

The authors declare that they have no competing interests.

FUNDING

The research source of funding is from non-tax revenue (PNBP) Faculty of Sport Science, State University of Malang, and researchers' self-funding.

ACKNOWLEDGMENTS

The authors of this study appreciate the Department of Sport Sciences, Faculty of Sport Sciences, State University of Malang and Physiology Laboratory, Faculty of Medicine Universitas Brawijaya Malang, Indonesia, for the provision of all samples employed and the laboratory to work for.

REFERENCES

- Adji, B.S., Sugiharto, Merawati, D., & Pranoto, A. (2021). The increase of uncoupling protein-1 expression after moderate intensity continuous exercises in obese females. *Jurnal SPORTIF : Jurnal Penelitian Pembelajaran*, 7(2), 194-205. https://doi.org/10.29407/js_unpgri.v7i2.15932.
- Aldiss P, Betts J, Sale C, Pope M, Budge H, Symonds ME. (2018). Exercise-induced 'browning' of adipose tissues. *Metabolism*. 81: 63-70. <https://doi.org/10.1016/j.metabol.2017.11.009>.
- Andarianto, A., Rejeki, P.S., Sakina, Pranoto, A., Seputra, T.W.A., Sugiharto, & Miftahussurur, M. (2022). Inflammatory markers in response to interval and continuous exercise in obese women. *Comparative Exercise Physiology*, 18(2), 135 – 142. <https://doi.org/10.3920/CEP210038>.
- Appari M, Channon KM, McNeill E. (2018). Metabolic Regulation of Adipose Tissue Macrophage Function in Obesity and Diabetes. *Antioxidants & Redox Signaling*. 29: 297-312. <https://doi.org/10.1089/ars.2017.7060>.
- Befroy DE, Petersen KF, Dufour S, Mason GF, Rothman DL, Shulman GI. (2008). Increased Substrate Oxidation and Mitochondrial Uncoupling in Skeletal Muscle of Endurance-Trained Individuals. *Proceedings of the National Academy of Sciences of the United States of America*. 105: 16701-16706.

- Brondani LA, Assmann TS, Duarte GCK, Gross JL, Canani LH, Crispim D. (2012). The role of the uncoupling protein 1 (UCP1) on the development of obesity and type 2 diabetes mellitus. *Arquivos Brasileiros de Endocrinologia & Metabologia*. 56: 215-225. <https://doi.org/10.1590/S0004-27302012000400001>.
- de Queiroz KB, Rodovalho GV, Guimarães JB, de Lima DC, Coimbra CC. (2021). Evangelista EA, Guerra-Sá R. Endurance training blocks uncoupling protein 1 up-regulation in brown adipose tissue while increasing uncoupling protein 3 in the muscle tissue of rats fed with a high-sugar diet. *Nutrition Research*. 32: 709-717. <https://doi.org/10.1016/j.nutres.2012.06.020>.
- Daskalopoulou SS, Cooke AB, Gomez Y-H, Mutter AF, Filippaios A, Mesfum ET, Mantzoros CS. (2014). Plasma irisin levels progressively increase in response to increasing exercise workloads in young, healthy, active subjects. *European Journal of Endocrinology*. 171(3): 343-352. <https://doi.org/10.1530/EJE-14-0204>.
- Dewal RS and Stanford KI. (2019). Effects of exercise on brown and beige adipocytes. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids*. 1864(1): 71-78. <https://doi.org/10.1016/j.bbalip.2018.04.013>.
- Dinas PC, Lahart IM, Timmons JA, Svensson PA, Koutedakis Y, Flouris AD, Metsios GS. (2017). Effects of physical activity on the link between PGC-1 α and FNDC5 in muscle, circulating Irisin and UCP1 of white adipocytes in humans: A systematic review. *F1000Research*. 6: 286. <https://doi.org/10.12688/f1000research.11107.1>.
- Fenzl, A. and Kiefer, F.W. (2014). Brown adipose tissue and thermogenesis. *Horm Mol Biol Clin Invest*. 19(1): 25–37. <https://doi.org/10.1515/hmbci-2014-0022>.
- Flouris AD, Dinas PC, Valente A, Andrade CMB, Kawashita NH, Sakellariou P. (2017). Exercise-induced effects on UCP1 expression in classical brown adipose tissue: a systematic review. *Hormone Molecular Biology and Clinical Investigation*. 31(2): 20160048. <https://doi.org/10.1515/hmbci-2016-0048>.
- Fuller-Jackson J-P and Henry BA. (2018). Adipose and skeletal muscle thermogenesis: studies from large animals. *Journal of Endocrinology*. 237: R99-R115. <https://doi.org/10.1530/JOE-18-0090>.
- Garneau L, Parsons SA, Smith SR, Mulvihill EE, Sparks LM, Aguer C. (2020). Plasma Myokine Concentrations After Acute Exercise in Non-obese and Obese Sedentary Women. *Frontiers in Physiology*. 11: 18. <https://doi.org/10.3389/fphys.2020.00018>.
- Gorski T, Mathes S, Krützfeldt J. (2018). Uncoupling protein 1 expression in adipocytes derived from skeletal muscle fibro/adipogenic

- progenitors is under genetic and hormonal control: UCP1 expression in skeletal muscle FAPs. *Journal of Cachexia, Sarcopenia and Muscle*. 9: 384-399. <https://doi.org/10.1002/jcsm.12277>.
- Herningtyas EH and Ng TS. (2019). Prevalence and distribution of metabolic syndrome and its components among provinces and ethnic groups in Indonesia. *BMC Public Health*. 19 (377). <https://doi.org/10.1186/s12889-019-6711-7>.
- Huh JY, Mougios V, Kabasakalis A, Fatouros I, Siopi A, Douroudos II, et al. (2014). Exercise-Induced Irisin Secretion Is Independent of Age or Fitness Level and Increased Irisin May Directly Modulate Muscle Metabolism Through AMPK Activation. *The Journal of Clinical Endocrinology & Metabolism*. 99(11): E2154–E2161. <https://doi.org/10.1210/jc.2014-1437>.
- Huh J.Y. (2018). The role of exercise-induced myokines in regulating metabolism. *Archives of Pharmacal Research*. 41: 14-29. <https://doi.org/10.1007/s12272-017-0994-y>.
- Jabbour G and Iancu H-D. (2017). High-intensity exercise training does not influence body weight but improves lipid oxidation in obese adults: a 6-week RCT. *BMJ Open Sport & Exercise Medicine*. 3(1): e000283. <https://doi.org/10.1136/bmjsem-2017-000283>.
- Khalafi M, Mohebbi H, Symonds ME, Karimi P, Akbari A, Tabari E, Faridnia M, Moghaddami K. (2020). The Impact of Moderate-Intensity Continuous or High-Intensity Interval Training on Adipogenesis and Browning of Subcutaneous Adipose Tissue in Obese Male Rats. *Nutrients*. 12: 925. <https://doi.org/10.3390/nu12040925>.
- Kim N, Kim J, Yoo C, Lim K, Akimoto T, Park J. (2018). Effect of acute mid-intensity treadmill exercise on the androgen hormone level and uncoupling protein-1 expression in brown fat tissue of mouse. *Journal of Exercise Nutrition & Biochemistry*. 22: 15-21. <https://doi.org/10.20463/jenb.2018.0003>.
- Kim SH and Plutzky J. (2016). Brown Fat and Browning for the Treatment of Obesity and Related Metabolic Disorders. *Diabetes & Metabolism Journal*. 40(1): 12-21. <https://doi.org/10.4093/dmj.2016.40.1.12>.
- Leal LG, Lopes MA, Batista ML. (2018). Physical Exercise-Induced Myokines and Muscle-Adipose Tissue Crosstalk: A Review of Current Knowledge and the Implications for Health and Metabolic Diseases. *Frontiers in Physiology*. 9: 1307. <https://doi.org/10.3389/fphys.2018.01307>.
- Mai S, Grugni G, Mele C, Vietti R, Vigna L, Sartorio A, Aimaretti G, Scacchi M, and Marzullo P. (2020). irisin levels in genetic and essential obesity: clues for a potential dual role. *Scientific Reports*. 10(1020): 1-9. <https://doi.org/10.1038/s41598-020-57855-5>.
- Moienneia N. and Hosseini S.R.A. (2016). Acute and chronic responses of metabolic myokine to different intensities of exercise in sedentary

young women. *Obesity Medicine*. 1: 15-20.
<http://dx.doi.org/10.1016/j.obmed.2015.12.002>.

Morrison S.F. (2016). Central control of body temperature. *F1000Research*. 5: F1000 Faculty Rev-880.
<https://doi.org/10.12688/f1000research.7958.1>.

Müller MJ, Enderle J, Bosity-Westphal A. (2016). Changes in Energy Expenditure with Weight Gain and Weight Loss in Humans. *Current Obesity Reports*. 5: 413-423. <https://doi.org/10.1007/s13679-016-0237-4>.

Oh K-J, Lee D, Kim W, Han B, Lee S, Bae K-H. (2017). Metabolic Adaptation in Obesity and Type II Diabetes: Myokines, Adipokines and Hepatokines. *International Journal of Molecular Sciences*. 18(1): 8. <https://doi.org/10.3390/ijms18010008>.

Paradisis GP, Zacharogiannis E, Mandila D, Smirtiotou A, Argeitaki P, Cooke CB. (2014). Multi-Stage 20-m Shuttle Run Fitness Test, Maximal Oxygen Uptake and Velocity at Maximal Oxygen Uptake. *Journal of Human Kinetics*.41: 81-87. <https://doi.org/10.2478/hukin-2014-0035>.

Purdom T, Kravitz L, Dokladny K, Mermier C. (2018). Understanding the factors that effect maximal fat oxidation. *J Int Soc Sports Nutr*.15 (3). <https://doi.org/10.1186/s12970-018-0207-1>.

Qiu S, Bosnyák E, Treff G, Steinacker JM, Nieß AM, Krüger K, Mooren FC, Zügel M, Schumann U. (2018). Acute exercise-induced irisin release in healthy adults: Associations with training status and exercise mode. *European Journal of Sport Science*.18: 1226-1233. <https://doi.org/10.1080/17461391.2018.1478452>.

Reisi, J., Ghaedi, K., Rajabi, H., & Marandi, S. M. (2016). Can Resistance Exercise Alter Irisin Levels and Expression Profiles of FNDC5 and UCP1 in Rats?. *Asian journal of sports medicine*. 7(4): e35205. <https://doi.org/10.5812/asjasm.35205>.

Rejeki, P. S., Baskara, P. G., Herawati, L., Pranoto, A., Setiawan, H. K., Lesmana, R., & Halim, S. (2022). Moderate-intensity exercise decreases the circulating level of betatrophin and its correlation among markers of obesity in women. *Journal of basic and clinical physiology and pharmacology*, 10.1515/jbcpp-2021-0393. Advance online publication. <https://doi.org/10.1515/jbcpp-2021-0393>.

Rejeki, P.S., Pranoto, A., Prasetya, R.E., & Sugiharto. (2021). Irisin serum increasing pattern is higher at moderate-intensity continuous exercise than at moderate-intensity interval exercise in obese females. *Comparative Exercise Physiology*, 17(5): 475–484. <https://doi.org/10.3920/CEP200050>.

Rodrigues KCC, Pereira RM, Campos TDP, Moura RF, Silva ASR, Cintra DE, Ropelle ER, Pauli JR, Araújo MB, Moura LP. (2018). The Role of Physical Exercise to Improve the Browning of White Adipose

- Tissue via POMC Neurons. *Frontiers in Cellular Neuroscience*. 12 (8). <https://doi.org/10.3389/fncel.2018.00088>.
- Sanchez-Delgado G, Martinez-Tellez B, Olza J, Aguilera CM, Gil Á, Ruiz JR. (2015). Role of Exercise in the Activation of Brown Adipose Tissue. *Ann Nutr Metab*. 67(1): 21-32. <https://doi.org/10.1159/000437173>.
- Shirvani, H. and Arabzadeh, E. (2020). Metabolic cross-talk between skeletal muscle and adipose tissue in high-intensity interval training vs. moderate-intensity continuous training by regulation of PGC-1 α . *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity*. 25:17–24. <https://doi.org/10.1007/s40519-018-0491-4>.
- Sugiharto, Merawati, D., Pranoto, A., Rejeki, P.S., Lupita, M.N., Adji, B.S., Susanto, H., & Taufiq, A. (2021). Acute Interval and Continuous Moderate-Intensity Exercise Enhanced Circadian Thermogenic Activity through Browning-related Genes in Obese Adolescent Female. *Malaysian Journal of Fundamental and Applied Sciences*, 17(5), 566-581. <https://doi.org/10.11113/mjfas.v17n5.2271>.
- Sugiharto, S., Merawati, D., Susanto, H., Pranoto, A., & Taufiq, A. (2022). The exercise-instrumental music program and irisin levels in younger non-professional athletes. *Comparative Exercise Physiology*, 18(1), 65–73. <https://doi.org/10.3920/CEP210015>.
- Sugiharto, Susanto, H, Andiana O, Merawati D. (2019a). Caloric Regulation Linked Thermogenesis in Acute Submaximal Intensity Exercise Model as The Effect of Audio Frequency Exposure. *IOP Conference Series: Materials Science and Engineering*. 515: 012069. <https://doi.org/10.1088/1757-899X/515/1/012069>.
- Sugiharto, Susanto H, Merawati D, Andiana O. (2019b). The Effect of Tempo of Musical Treatment and Acute Exercise on Vascular Tension and Cardiovascular Performance: A Case Study on Trained Non-Athletes. *IOP Conference Series: Materials Science and Engineering*. 515: 012033. <https://doi.org/10.1088/1757-899X/515/1/012033>.
- Tsuchiya Y, Ando D, Goto K, Kiuchi M, Yamakita M, Koyama K. (2014). High-Intensity Exercise Causes Greater Irisin Response Compared with Low-Intensity Exercise under Similar Energy Consumption. *The Tohoku Journal of Experimental Medicine*. 233(2): 135-140. <https://doi.org/10.1620/tjem.233.135>.
- Tsuchiya Y, Ando D, Takamatsu K, Goto K. (2015). Resistance exercise induces a greater irisin response than endurance exercise. *Metabolism: Clinical and Experimental*. 64(9): 1042-1050. <https://doi.org/10.1016/j.metabol.2015.05.010>.
- Zhang J, Valverde P, Zhu X, Murray D, Wu Y, Yu L, Jiang H, Dard MM, Huang J, Xu Z, Tu Q, Chen J. (2017). Exercise-induced irisin in bone

and systemic irisin administration reveal new regulatory mechanisms of bone metabolism. *Bone Research*. 5: 1-14. <https://doi.org/10.1038/boneres.2016.56>.